

University of Groningen

Epigenetic reprogramming of endogenous genes for permanent modulation of gene expression

Huisman, Christian

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version

Publisher's PDF, also known as Version of record

Publication date:

2015

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Huisman, C. (2015). *Epigenetic reprogramming of endogenous genes for permanent modulation of gene expression: Targeted interventions by self-engineered zinc finger proteins*. [Thesis fully internal (DIV), University of Groningen]. University of Groningen.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.



General introduction and Scope of the thesis

Chapter 1

General introduction

The human body is made up of more than 200 different cell types and each cell type has its own specific function. The genetic information, encoded in the DNA, instructs the cells how to perform its function. This genetic information contains roughly 20700 genes (1), which code for the functional units of the cell; the proteins. The specific protein expression pattern displayed by a cell determines which biological processes are taken place. Tight regulation of these processes, including growth, differentiation and proliferation ensure normal cell functioning. Usually only a subset of the genes is transcribed to proteins in order for the cell to perform its function.

The number of genes that are actively transcribed in a cell is constantly changing, and dynamically regulated by a layer on top of the DNA called the 'epigenetic layer'. Epigenetics is the additional layer of heritable information on top of the base pair sequence of the genome (2). The structural units of the epigenetic layer are the nucleosomes (3), which are composed of several types of histone proteins. Histone proteins are highly tunable through their protruding tails which can be posttranslational modified on many positions. The nucleosomes organize and package the DNA into the nucleus of a cell into highly ordered structures allowing for flexible modulation of gene expression patterns.

The epigenetic state of a gene is an important factor that determines the expression level of a gene (4). The two main components of epigenetic regulation (Figure 1) are DNA methylation; the addition of methyl group on the 5th position of the cytosine ring within CpG dinucleotides (CpGs) (5), and posttranslational histone modifications (4). A variety of histone modifications have been identified, including histone methylation (e.g. trimethylation of lysine 9 of histone 3) and histone (de)acetylation (e.g. (de)acetylation of histone 3). Methylation of DNA and histones is usually associated with low levels of gene expression, while unmethylated DNA and histone acetylation is usually associated with high levels of gene expression (4). However, there are several exceptions, for example the very well-studied histone mark trimethylation of lysine 4 of histone H3 is usually associated with high levels of transcription (4). Especially epigenetic modifications found in the promoter region of a gene (region of DNA that initiates transcription) determine if a gene is transcribed or not. In 40% of the genes, promoter regions contain CpG islands (DNA sequences with a high frequency of CpGs) (6). The DNA methylation levels of these CpG islands often determine if the gene is 'off' or 'on', while the histone modifications may play a role in determining the precise levels of gene expression.

Epigenetics in cancer

It becomes increasingly clear that many diseases are associated with epigenetic alterations causing aberrant gene expression (7). A common example is cancer.




Figure 1. DNA methylation and histone modifications. Schematic representation of the epigenetic layer on top of the DNA sequence which influence gene expression.

Every year, more than 10 million people die from this disease, and the number of new cases is increasing rapidly. Therefore, cancer is one of the leading causes of death. Cancer is characterized by uncontrolled cell growth, as genes that stimulate cell proliferation (oncogenes) are overexpressed or gained function, while genes inhibiting these processes (tumor suppressor genes (TSGs)) are downregulated or dysfunctional. Although cancer has long been considered as a set of diseases caused by genetic abnormalities, new insights after intensive research justify the concept that cancer is an epigenetic disease as well. Epigenetic deregulation can lead to genomic instability and influences the gene expression pattern displayed by cells. The epigenetic dysregulation may even be the key that drives carcinogenesis (8). Besides that, most cancers harbor inactivating mutations in genes that control the epigenome, which further contribute to epigenetic deregulation and cancer progression (8). The most observed epigenetic abnormality in malignant cells is promoter DNA hypermethylation of CpG islands, which in turn are accompanied by the presence of repressive histone marks and a loss of activating histone marks, which further inhibit gene expression (9). For example, in various types of cancer, *Maspin*, *EPB41L3*, *CDKN2A* and *RASSF1A* are among the genes which are down-regulated by epigenetic mechanisms (10, 11) and are subject of investigation in the current thesis.

Cervical cancer is one of cancers which is extensively studied in this thesis, and is one of the top killers of women by cancer worldwide. In the last decade, genome-wide mapping of DNA methylation in normal- and cervical cancer

genomes revealed that a large number of genes in cervical cancer are associated with hypermethylated gene promoters (12), (11). Besides their functional implications of gene inactivation in tumor development, these methylation patterns can serve as diagnostic biomarkers for cervical cancer. Current screenings programs rely on the pap-smear assays which have poor sensitivity. Our group and others have reported several gene promoters that are specifically hypermethylated in cervical cancer. One of the hypermethylation markers we identified was *C13ORF18* (10), but we and others also found that *CCNA1*, *EPB41L3* and *TFPI2* methylation levels were strongly associated with cancer (11, 13-17). As promoter hypermethylation is often seen for TSGs, such potent diagnostic marker genes might exhibit tumor suppressive activities upon re-expression. Therefore, one of the goals in this thesis was to re-express these silenced marker genes and study their tumor suppressive activity in cervical cancer cell lines. Key TSGs for cervical cancer growth may subsequently be used as therapeutic target to decrease tumor growth in this malignancy.

It has already been shown that re-introduction of single genes into cancer cells where the gene is inactivated showed potent anti-tumor effects, as demonstrated for *CDKN2A* (encoding p16Ink4A) (18) or *TP53* (19). Contrary to genes that are inactivated through genetic mutations, epigenetic deregulation is a reversible process and can be targeted by epigenetic drugs (20). Current clinical drugs, such as the DNA methylation inhibitor Decitabin (5-aza-deoxycytosine) and the histone deacetylase inhibitor Vorinostat (suberoylanilide hydroxamic acid, SAHA), induce re-expression of a variety of hypermethylated silenced genes (21). So far, four epigenetic drugs have been FDA approved for subtypes of leukemias and lymphomas, and ongoing clinical trials in solid tumors indicate some beneficial effects for these patients as well (22). However, the disadvantages of these drugs include their genome-wide effects (20). In that respect it is important to note that epigenetic drugs have been shown to drive tumor promotion and metastasis by enhanced expression of pro-metastatic genes (23). Therefore, gene-targeted re-activation of TSGs is of great importance. A novel way to achieve gene-targeted re-activations is by the use of artificial transcription factors (ATFs) (24).

Artificial transcription factors (ATFs)

An interesting development in the last years for the gene-targeted re-expression of silenced genes or the downregulation of overexpressed genes is the use of ATFs (24). ATFs consist of a transcriptional regulatory domain coupled to an engineered DNA binding domain (Figure 2), such as a Zinc Finger Protein (ZFP), a transcription activator-like effector (TALE) protein or a Clustered Regularly Interspaced Short Palindromic Repeats (CRISPR) associated 9 (Cas9) protein. Such DNA binding domains can be designed to target virtually any gene in the human genome (25, 26). While the TALE and CRISPR/CAS9 system are relative new targeting platforms, ZFP-mediated targeting has already been explored for more than a decade, and

this system is best characterized for its ability to target and modulate gene expression. ZFPs are natural occurring transcription factors containing several DNA binding domains (“fingers”). Each finger is around 30 amino acids and recognizes three base pairs. Individual fingers can be stitched together to increase their specificity. Typically, ZFPs are designed to target a unique DNA stretch of 18 base pairs within the genomic DNA. An amino acid- DNA binding code has been generated allowing ZFPs to target the desired triplet (27). The most common transcriptional regulatory domains are the gene-activator VP64 (a tetramer of the herpes simplex virus gene activator VP16) and the gene-repressor SKD




Figure 2. Artificial transcription factors (ATFs). ATFs are fusion proteins consisting of a DNA binding domain (DBD), e.g. an engineered Zinc Finger Protein (ZFP), fused to a transcriptional effector domain. The DBD is targeted to the gene of interest. The effector domain enforces the modulation of gene expression (VP64, SKD). The approach can be redesigned to directly modulate the epigenetic landscape (Epigenetic Editing) depending on which effector domain (e.g. Tet2, G9a, SUV39H1 and M.Sss1) is used.

(Krüppel associated box repressor domain). ATFs technology has proven itself as very useful tool for the modulation of gene expression in many disease models (24) and ZFP-targeting is currently the most advanced DNA targeting approach. This technology has the potency to activate gene expression, but depending on the transcriptional regulatory domain, can also down-regulate gene-expression (28). Compared to the current drugs which target only 2.5% of the proteins, the ability to target any gene using ATFs, including previously undruggable targets (29, 30), is a huge advantage. Traditional cDNA expression approaches have the disadvantage that only one splice variant is expressed, but ATFs enable expression from the endogenous gene locus allowing for all splice variants to be expressed in natural ratios. The importance of expressing alternative splice variants have been demonstrated for the *VEGF* gene (31) and was the basis for the first clinical trial based on ATFs. Furthermore, ATFs achieve efficient gene silencing by targeting the

DNA molecule directly, thereby overcoming the difficulties of RNA interference which needs to target continuously the newly produced mRNA molecules. The above mentioned advantages make ATFs suitable to study gene function or to modulate gene expression for therapeutic purposes. Interestingly, the ZFPs can also be attached to nucleases for site-specific DNA editing (changing of the genetic sequence at a specific site). Genome editing with engineered nucleases was declared 'Method of the Year 2011' by Nature Methods (32) and currently several clinical phase II ZF-nuclease trials are ongoing.

Epigenetic editing

The transcriptional effects of VP64 and SKD are traditionally considered transient in differentiated cells, as no epigenetic marks are actively rewritten, and genes likely return to the 'normal' state after the ATF is removed (33, 34). For 'permanent' modulation of gene expression, the epigenetic marks need to be rewritten, which may be achieved by a technique called *Epigenetic Editing* (35). Epigenetic editing is the gene-specific targeting of epigenetic enzymes to overwrite epigenetic signatures and to achieve long-term a) silencing of actively transcribed genes or b) re-expression of epigenetically silenced genes. Targeted epigenetic enzymes have the potency to change local DNA methylation or histone modifications at the site, which may result in transcriptional modulation. Epigenetic editing offers attractive prospects as the introduced epigenetic changes may achieve stable reprogramming, even after clearance of the drug (hit and run approach). So far, this exciting novel strategy has already resulted in the efficient transcriptional modulation of various genes targeting either histone modifications or DNA methylation. *VEGF* was the first gene modified by epigenetic editing; the catalytic domains of histone methyl transferases G9a and SUV39-H1 were targeted to the *VEGF* promoter, resulting in efficient downregulation by the induced repressive histone marks (36). The subsequent targeting of methyltransferase (DNA methylating enzyme) to the promoter regions of *Sox2*, *Maspin* and *VEGF* demonstrated that downregulation can also be achieved by targeted DNA methylation (37, 38). In this thesis, we further explored such targeting of histone modifying enzymes and DNA methylating enzymes for their repressive effects on gene expression.

To achieve long term re-expression of epigenetically silenced genes by means of epigenetic editing, epigenetically activating enzymes or enzymes that remove repressive marks, such as DNA methylation, can be targeted to silenced promoters. Recently, the ten eleven translocation (Tet) enzymes have been identified as actively DNA demethylators (39). Therefore, Tet proteins can be fused to ZFPs to achieve site-specific demethylation potentially resulting in gene-activation. For a long time, gene activation by targeted DNA demethylation seemed unfeasible in mammalian cells as no enzymes were identified with the capacity to actively demethylate DNA, although various biological processes

hinted to the existence of such mechanism, including the rapid demethylation of the paternal genome in the zygote hours after fertilisation (40). With the identification of the Tet-enzymes and other enzymes implicated in the DNA demethylating pathway, the feasibility of targeted DNA demethylation to activate gene expression is under exploration and receives widespread attention (41). Recently, the first studies appeared which achieved demethylation and upregulation of genes by targeting DNA demethylases to hypermethylated promoters (42, 43). Also in this thesis, we investigated the ability of Tet-enzymes (Tet2) to demethylate and activate hypermethylated genes.

Scope of the thesis

This thesis describes how ZFP-based ATF-technology can be applied to unravel a biological function of silenced genes currently arising from biomarker searches in cervical cancer and other malignancies. Furthermore, this thesis aims to provide more insights into causal effects of epigenetic marks on gene expression patterns (DNA methylation, histone modifications) and crosstalk in-between. Moreover, we explored the potencies of epigenetic editing to achieve stable expression modulation and we contribute to unravelling the underlying epigenetic rules for such epigenetic reprogramming. Current approaches for stable re-expression modulation require the potentially hazardous integration of viral DNA into the host genome, which limit their clinical use. The presented hit-and-run strategy circumvents these integrations, and assists to realize the curable genome concept, allowing a major role in modulation of expression of any gene at will, including currently undruggable (non)protein-coding genes.

In total, we utilized 38 ZFPS (34 self-engineered ZFPs) targeted to eight different genes (Table I), fused them to transcriptional expression modulators (VP64, SKD) or epigenetic enzymes (Tet2, G9A, SUV39H1 and M.Sss1) and studied the effect on various cell biological features, including target gene expression modulation, cell growth, DNA methylation and histone modifications.

Targeting the EpCAM and ICAM-1 promoter to decrease tumor growth in ovarian cancer

The first gene we modulated by ATF-technology was our model gene epithelial cell adhesion molecule (*EpCAM*), which is overexpressed in most carcinomas. Dependent on the tumor type, its overexpression is either associated with improved or worse patient survival. For ovarian cancer however, the role of EpCAM remains unclear. We used previously engineered ATFs for both the up- and downregulation of EpCAM. The bidirectional ATF-based approach is uniquely suited to study cell-type-specific biological effects of EpCAM expression. Using this approach, the function of EpCAM in ovarian cancer was investigated (**Chapter 2**).

Table I. The gene-targeted ATFs used in this thesis.

Gene	Function	ATFs
<i>EPCAM</i>	Calcium-independent cell adhesion molecule overexpressed on cancer cells	2
<i>ICAM-1</i>	Cell adhesion molecule involved in immune response (44)	1*
<i>C13ORF18</i>	Putative TSG with unknown function, possibly involved in autophagy regulation	5
<i>CCNA1</i>	Cyclin involved in control of (germ line meiotic) cell cycle	6
<i>TFPI2</i>	Linked to regulation of plasmin-mediated matrix remodeling	11
<i>Maspin</i>	Inhibits cell motility and invasion, inhibitor of HDAC-1	2*
<i>EPB41L3</i>	Cell adhesion, cell motility and cell growth	2
<i>HER2</i>	Enhances activation of signaling pathways promoting cell proliferation (45)	1*
<i>PLOD2</i>	Catalyses the hydroxylation of lysyl residues in collagen-like peptides	8

*from literature

In the next chapter (**Chapter 3**), we investigated whether ICAM-1 can function as TSG in ovarian cancer in the absence of immune cells. ICAM-1 is a protein which is commonly expressed on endothelial cells and cells of the immune system. In ovarian cancer, ICAM-1 is downregulated compared with healthy ovarian cells and expression is correlated with decreased tumorigenicity (46). Whereas ICAM-1 expression on tumor cells is of importance for attracting immune cells, ICAM-1 might also induce tumorigenicity and chemoresistance. In ovarian cancer, such a role of ICAM-1 is unclear. We investigated whether ICAM-1 has a tumor suppressive role in this malignancy by re-expression of ICAM-1 using previously published *ICAM-1*-targeting ATFs (44).

Re-expression of TSGs in cervical cancer

To investigate whether we could achieve re-expression of genes which were aberrantly silenced by promoter hypermethylation in cervical cancer, we performed three different studies. In the first study, we aimed to re-express *C13ORF18* in cervical cancer cell lines by five engineered ATFs and studied the effect of re-expression on cancer-related cell biological features. Currently, the function of *C13ORF18* is unknown, but this gene is specifically methylated in cervical cancer compared to normal cervix (10). This supposed silenced state of *C13ORF18* in cervical cancer hints towards a tumor suppressive role for this gene

in this malignancy, which was elaborately explored in this chapter. Moreover, in this chapter we describe the effects of targeting ATFs to the *C13ORF18* promoter on the local DNA methylation levels and histone modifications (**Chapter 4**).

In the second study (**Chapter 5**), we aimed to re-activate the hypermethylated candidate tumor suppressor genes *C13ORF18*, *CCNA1*, *TFPI2* and *Maspin* by Tet-induced demethylation. To target Tet2 to hypermethylated CpGs, we used engineered zinc fingers proteins (ZFPs), which were all validated for gene regulatory capacity by fusions to VP64. We also compared the hypothesized tumor suppressive effects of these genes by the engineered gene-targeting constructs as well as by cDNA overexpression. We used primary spherical growing cell lines to study the functional effects, which are a better representative compared to established cell lines, and may gap the bridge with real tumors. Furthermore, we addressed the binding of ZFPs to their intended sequence and throughout the genome using ChIP-Seq and elaborate on the 'questionable' role of *Maspin* as TSG (47) in (cervical) cancer.

In the third study, we aimed to fully reprogram the biomarker *EPB41L3* in order to improve the sustainability of gene re-expression using the synergistic effects of ATFs with epigenetic drugs (5-aza-dC or TSA). Until now, no methods exist to permanently re-activate single silenced genes, and this chapter describes a novel way to improve the sustainability of targeted re-expression. We also addressed the potential synergy of ATFs with epigenetic drugs. Interestingly, *EPB41L3* was recently identified as the best marker for detecting CIN2/3 (13, 17), but its function in cervical cancer has not been explored yet (**Chapter 6**).

Downregulation of gene-expression by epigenetic editing

In order to down-regulate gene expression, we targeted Human Epidermal Growth Factor Receptor 2 (*HER2*) linked to repressive epigenetic editing domains. *HER2* is an important member of the epidermal growth factor receptor family and overexpressed in several cancer types. Downregulation of this oncogene dramatically decreases cancer cell metabolism, growth and proliferation. In this chapter, we targeted the *HER2* promoter using a previously described ZFP specific for the *HER2* promoter (45) and fused it to histone methyl transferases G9a and SUV39H1. Subsequently, we investigated if we could overwrite the histone modifications in the *HER2* promoter and whether this results in *HER2* expression modulation and reduced tumorigenicity (**Chapter 7**).

To further exploit the potentials of ATFs and *Epigenetic Editing* for the downregulation of gene expression, we set out to target procollagen-lysine, 2-oxoglutarate 5-dioxygenase 2 (*PLOD2*). Excessive collagen deposition is the hallmark of fibrosis (48), and downregulation of *PLOD2* is believed to inhibit this process. For the silencing of target genes, both targeted methylation of DNA and histones has been explored. However, insights into sustainability of gene expression modulation by such methods are largely lacking. In this chapter, we

designed a system to compare the effects of a strong DNA methylator (M.SssI) and a strong gene repressor (SKD) on sustainability of gene repression of *PLOD2*. We engineered stable inducible human skin fibroblasts to express ZFP-SKD or ZFP-M.SssI fusions targeting the *PLOD2* promoter. After short term expression of the *PLOD2*-targeting constructs, we investigated the effects on gene expression and local epigenetic features (DNA methylation and histone marks). Finally, we aimed to downregulate *PLOD2* expression in a profibrotic environment (mimicked by addition of TFGb) to demonstrate the therapeutic potential of our tools (**Chapter 8**). In **Chapter 9** the results from chapter 2-8 are summarized and discussed.

References

1. Harrow J, Frankish A, Gonzalez JM, et al. GENCODE: The reference human genome annotation for the ENCODE project. *Genome Res* 2012;22(9):1760-74.
2. Park YJ, Claus R, Weichenhan D, Plass C. Genome-wide epigenetic modifications in cancer. *Prog Drug Res* 2011;67:25-49.
3. Van Holde KE, Sahasrabuddhe CG, Shaw BR. A model for particulate structure in chromatin. *Nucleic Acids Res* 1974;1(11):1579-86.
4. ENCODE Project Consortium, Bernstein BE, Birney E, et al. An integrated encyclopedia of DNA elements in the human genome. *Nature* 2012;489(7414):57-74.
5. Holliday R, Pugh JE. DNA modification mechanisms and gene activity during development. *Science* 1975;187(4173):226-32.
6. van Vlerken LE, Hurt EM, Hollingsworth RE. The role of epigenetic regulation in stem cell and cancer biology. *J Mol Med (Berl)* 2012;90(7):791-801.
7. Baylin SB, Jones PA. A decade of exploring the cancer epigenome - biological and translational implications. *Nat Rev Cancer* 2011;11(10):726-34.
8. You JS, Jones PA. Cancer genetics and epigenetics: Two sides of the same coin? *Cancer Cell* 2012;22(1):9-20.
9. Jones PA. Functions of DNA methylation: Islands, start sites, gene bodies and beyond. *Nat Rev Genet* 2012;13(7):484-92.
10. Yang N, Eijssink JJ, Lendvai A, et al. Methylation markers for CCNA1 and C13ORF18 are strongly associated with high-grade cervical intraepithelial neoplasia and cervical cancer in cervical scrapings. *Cancer Epidemiol Biomarkers Prev* 2009;18(11):3000-7.
11. Hoque MO, Kim MS, Ostrow KL, et al. Genome-wide promoter analysis uncovers portions of the cancer methylome. *Cancer Res* 2008;68(8):2661-70.
12. Lendvai A, Johannes F, Grimm C, et al. Genome-wide methylation profiling identifies hypermethylated biomarkers in high-grade cervical intraepithelial neoplasia. *Epigenetics* 2012;7(11):1268-78.
13. Vasiljevic N, Scibior-Bentkowska D, Brentnall AR, Cuzick J, Lorincz AT. Credentialing of DNA methylation assays for human genes as diagnostic biomarkers of cervical intraepithelial neoplasia in high-risk HPV positive women. *Gynecol Oncol* 2014.
14. Sova P, Feng Q, Geiss G, et al. Discovery of novel methylation biomarkers in cervical carcinoma by global demethylation and microarray analysis. *Cancer Epidemiol Biomarkers Prev* 2006;15(1):114-23.
15. Santin AD, Zhan F, Bignotti E, et al. Gene expression profiles of primary HPV16- and HPV18-infected early stage cervical cancers and normal cervical epithelium: Identification of novel candidate molecular markers for cervical cancer diagnosis and therapy. *Virology* 2005;331(2):269-91.

16. Kitkumthorn N, Yanatatsanajit P, Kiatpongsan S, et al. Cyclin A1 promoter hypermethylation in human papillomavirus-associated cervical cancer. *BMC Cancer* 2006;6:55.
17. Eijssink JJ, Lendvai A, Deregowski V, et al. A four-gene methylation marker panel as triage test in high-risk human papillomavirus positive patients. *Int J Cancer* 2012;130(8):1861-9.
18. Lu Y, Zhang X, Zhang J. Inhibition of breast tumor cell growth by ectopic expression of p16/INK4A via combined effects of cell cycle arrest, senescence and apoptotic induction, and angiogenesis inhibition. *J Cancer* 2012;3:333-44.
19. Hamada K, Alemany R, Zhang WW, et al. Adenovirus-mediated transfer of a wild-type p53 gene and induction of apoptosis in cervical cancer. *Cancer Res* 1996;56(13):3047-54.
20. Rius M, Lyko F. Epigenetic cancer therapy: Rationales, targets and drugs. *Oncogene* 2012;31(39):4257-65.
21. Rodriguez-Paredes M, Esteller M. Cancer epigenetics reaches mainstream oncology. *Nat Med* 2011;17(3):330-9.
22. Juergens RA, Wrangle J, Vendetti FP, et al. Combination epigenetic therapy has efficacy in patients with refractory advanced non-small cell lung cancer. *Cancer Discov* 2011;1(7):598-607.
23. Yu Y, Zeng P, Xiong J, Liu Z, Berger SL, Merlino G. Epigenetic drugs can stimulate metastasis through enhanced expression of the pro-metastatic ezrin gene. *PLoS One* 2010;5(9):e12710.
24. Sera T. Zinc-finger-based artificial transcription factors and their applications. *Adv Drug Deliv Rev* 2009;61(7-8):513-26.
25. Uil TG, Haisma HJ, Rots MG. Therapeutic modulation of endogenous gene function by agents with designed DNA-sequence specificities. *Nucleic Acids Res* 2003;31(21):6064-78.
26. Gersbach CA, Perez-Pinera P. Activating human genes with zinc finger proteins, transcription activator-like effectors and CRISPR/Cas9 for gene therapy and regenerative medicine. *Expert Opin Ther Targets* 2014;18(8):835-9.
27. Carroll D, Morton JJ, Beumer KJ, Segal DJ. Design, construction and in vitro testing of zinc finger nucleases. *Nat Protoc* 2006;1(3):1329-41.
28. Gersbach CA, Gaj T, Barbas CF, 3rd. Synthetic zinc finger proteins: The advent of targeted gene regulation and genome modification technologies. *Acc Chem Res* 2014;47(8):2309-18.
29. Stolzenburg S, Rots MG, Beltran AS, et al. Targeted silencing of the oncogenic transcription factor SOX2 in breast cancer. *Nucleic Acids Res* 2012;40(14):6725-40.
30. Ji Q, Fischer AL, Brown CR, et al. Engineered zinc-finger transcription factors activate OCT4 (POU5F1), SOX2, KLF4, c-MYC (MYC) and miR302/367. *Nucleic Acids Res* 2014;42(10):6158-67.

31. Rebar EJ, Huang Y, Hickey R, et al. Induction of angiogenesis in a mouse model using engineered transcription factors. *Nat Med* 2002;8(12):1427-32.
32. Baker M. Gene-editing nucleases. *Nat Methods* 2012;9(1):23-6.
33. Beltran AS, Russo A, Lara H, Fan C, Lizardi PM, Blancafot P. Suppression of breast tumor growth and metastasis by an engineered transcription factor. *PLoS One* 2011;6(9):e24595.
34. Tan S, Guschin D, Davalos A, et al. Zinc-finger protein-targeted gene regulation: Genomewide single-gene specificity. *Proc Natl Acad Sci U S A* 2003;100(21):11997-2002.
35. de Groote ML, Verschure PJ, Rots MG. Epigenetic editing: Targeted rewriting of epigenetic marks to modulate expression of selected target genes. *Nucleic Acids Res* 2012;40(21):10596-613.
36. Snowden AW, Gregory PD, Case CC, Pabo CO. Gene-specific targeting of H3K9 methylation is sufficient for initiating repression in vivo. *Curr Biol* 2002;12(24):2159-66.
37. Rivenbark AG, Stolzenburg S, Beltran AS, et al. Epigenetic reprogramming of cancer cells via targeted DNA methylation. *Epigenetics* 2012;7(4).
38. Siddique AN, Nunna S, Rajavelu A, et al. Targeted methylation and gene silencing of VEGF-A in human cells by using a designed Dnmt3a-Dnmt3L single-chain fusion protein with increased DNA methylation activity. *J Mol Biol* 2013;425(3):479-91.
39. Tahiliani M, Koh KP, Shen Y, et al. Conversion of 5-methylcytosine to 5-hydroxymethylcytosine in mammalian DNA by MLL partner TET1. *Science* 2009;324(5929):930-5.
40. Oswald J, Engemann S, Lane N, et al. Active demethylation of the paternal genome in the mouse zygote. *Curr Biol* 2000;10(8):475-8.
41. Voigt P, Reinberg D. Epigenome editing. *Nat Biotechnol* 2013;31(12):1097-9.
42. Maeder ML, Angstman JF, Richardson ME, et al. Targeted DNA demethylation and activation of endogenous genes using programmable TALE-TET1 fusion proteins. *Nat Biotechnol* 2013;31(12):1137-42.
43. Chen H, Kazemier HG, de Groote ML, Ruiters MH, Xu GL, Rots MG. Induced DNA demethylation by targeting ten-eleven translocation 2 to the human ICAM-1 promoter. *Nucleic Acids Res* 2014;42(3):1563-74.
44. Magnenat L, Blancafot P, Barbas CF,3rd. In vivo selection of combinatorial libraries and designed affinity maturation of polydactyl zinc finger transcription factors for ICAM-1 provides new insights into gene regulation. *J Mol Biol* 2004;341(3):635-49.
45. Beerli RR, Dreier B, Barbas CF,3rd. Positive and negative regulation of endogenous genes by designed transcription factors. *Proc Natl Acad Sci U S A* 2000;97(4):1495-500.
46. Arnold JM, Cummings M, Purdie D, Chenevix-Trench G. Reduced expression of intercellular adhesion molecule-1 in ovarian adenocarcinomas. *Br J Cancer* 2001;85(9):1351-8.

47. Teoh SS, Vieusseux J, Prakash M, et al. Maspin is not required for embryonic development or tumour suppression. *Nat Commun* 2014;5:3164.
48. van der Slot AJ, Zuurmond AM, Bardoel AF, et al. Identification of PLOD2 as telopeptide lysyl hydroxylase, an important enzyme in fibrosis. *J Biol Chem* 2003;278(42):40967-72.